

## CAUTION

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Nufarm  
**DRIVE**<sup>®</sup>  
Herbicide

**ACTIVE CONSTITUENT: 750 g/kg QUINCLORAC**

GROUP II HERBICIDE

For the post-emergence control of summer grass and white clover and the suppression of Kikuyu in turf as specified in the DIRECTIONS FOR USE table.

Contents: 2 kg

Nufarm Australia Limited  
ACN 004 377 780  
103-105 Pipe Road  
Laverton North Victoria 3026  
Tel: (03) 9282 1000  
Fax: (03) 9282 1001



® = Registered trademark of BASF

**DIRECTIONS FOR USE:****RESTRAINTS:**

Clippings from the first three mowings should be left on the treated area.

DO NOT apply into any ornamental bed

DO NOT apply when weeds are not actively growing

DO NOT apply if heavy rain is expected within 48 hours, or within 8 hours prior to irrigation.

DO NOT apply this product through any type of irrigation system

DO NOT apply to golf course collars or greens

DO NOT apply within 4 weeks after seedling emergence of Kentucky Bluegrass, Creeping Bentgrass and Perennial Ryegrass.

| SITUATION   | WEEDS CONTROLLED  | Rate kg/ha         | Critical Comments  |
|---|---|--------------------|--|
| <b>Established turf of:</b><br>Common couch<br><i>(Cynodon dactylon)</i> ,<br>hybrid couch<br><i>(Cynodon dactylon x C. transvaalensis)</i> ,<br>Japanese lawngrass<br><i>(Zoysia japonica)</i> ,<br>marine couch<br><i>(Sporobolus virginicus)</i> ,<br>perennial ryegrass<br><i>(Lolium perenne)</i><br>bent<br><i>(Agrostis stolonifera)</i> ,<br>Kentucky Bluegrass<br><i>(Poa pratensis)</i> | Summer grass<br><i>(Digitaria spp.)</i>                       | 1.1 in 400 L water | <b>One application:</b><br>Weeds at cotyledon stage up to prior to second tiller<br>OR<br>Weeds at 5 tillers and greater<br><br><b>Two applications: 21 to 28 days apart:</b><br>Weeds from 2 to 4 tillers<br><br>The addition of a crop oil adjuvant may improve weed control in unfavourable weather conditions. |
|   | Kikuyu <i>(Pennisetum clandestinum)</i><br>(Suppression only) |                    | For Kikuyu suppression, apply twice, 14 to 21 days apart. Best results are gained from Autumn applications.<br><br>The addition of a crop oil adjuvant may improve weed control in unfavourable weather conditions.  |
| See also<br>"Protection of crops,<br>native and other non-<br>target plants" section  | White Clover<br><i>(Trifolium repens)</i>                     |                    | Apply to actively growing weeds.   |

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

## GENERAL INSTRUCTIONS

DRIVE will provide post-emergence control of summer grass and clover in turf. Quinclorac is rapidly absorbed through the foliage and translocated throughout the plant. Death of sensitive weeds usually occurs between 1 and 2 weeks after treatment. Kikuyu can be suppressed by applying two post-emergence treatments. Best results will be achieved if Kikuyu is treated in Autumn.

## HERBICIDE RESISTANCE WARNING

### GROUP I HERBICIDE

Nufarm DRIVE Herbicide is a member of the Carboxylic acid group of herbicides and has the Disruptors of Plant Cell Growth mode of action. For weed resistance management this product is a Group I herbicide. Some naturally occurring weed biotypes resistant to DRIVE and other disruptors of plant cell growth herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. The resistant weeds will not be controlled by DRIVE or other Group I herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Nufarm Australia Limited accepts no liability for any losses that may result from the failure of DRIVE to control resistant weeds.

## MIXING

To ensure even mixing, half-fill the spray tank with clean water and add the required amount of product. Agitate thoroughly, then add the remainder of the water. Agitate again before spraying commences.

The addition of a crop oil adjuvant may improve weed control in unfavourable weather conditions.

## APPLICATION

Apply DRIVE to actively growing weeds at the growth stage indicated in the Directions for Use table. Apply the product in 400 L of water/ha. DO NOT apply through any type of irrigation system. Avoid overlaps during application. Separate applications should be made if all target weeds are not at the correct growth stage for treatment at the same time. It is recommended not to mow 2 days before or after applying DRIVE to maximize weed control and minimize turf injury. Refer to the Restraints listed above the Directions for Use table for further application advice.

## COMPATIBILITY

The product is compatible with DSMA, Primo\*250EC and Primo\*Maxx.

## PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift

onto nearby susceptible plants/crops, cropping lands, pastures or native vegetation.

Quinclorac has properties and characteristics associated with chemicals detected in groundwater. The use of this product where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

DO NOT apply to Bahia grass (*Paspalum notatum*), centipedegrass (*Eremochloa ophiuroides*), buffalo grass (*Stenotaphrum secundatum*), Kikuyu (*Pennisetum clandestinum*), Qld blue Couch (*Digitaria didactyla*), carpet grass (*Axonopus* spp.), or lawns or turf where desirable clovers are present.

DO NOT apply to ornamental beds or to exposed feeder roots of trees and ornamentals. Be particularly careful within the drip line of these plants. DO NOT apply to golf course collars or greens.

Clippings from the first three mowings should be left on the treated area.

## PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

## STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If not available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should NOT be burnt.

## FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

## SAFETY DIRECTIONS

Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container, preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and a washable hat, and elbow length PVC gloves. Wash hands after use. After each day's use wash gloves and contaminated clothing.

## MATERIAL SAFETY DATA SHEET (MSDS)

For further information refer to the Material Safety Data Sheet (MSDS).

**CONDITIONS OF SALE**

“Any provisions or rights under the Trade Practices Act 1974 or relevant state legislation which cannot be excluded by those statutes or by law are not intended to be excluded by these conditions of sale. Subject to the foregoing, all warranties, conditions, rights and remedies, expressed or implied under common law, statute or otherwise, in relation to the sale, supply, use or application of this product, are excluded. Nufarm Australia Limited and/or its affiliates (“Nufarm”) shall not accept any liability whatsoever (including consequential loss), or howsoever arising (including negligence) for any damage, injury or death connected with the sale, supply, use or application of this product except for liability which cannot be excluded by statute.”

**In case of emergency: Phone 1800 033 498**

Ask for shift supervisor. Toll free 24 hours.

APVMA Approval No.: 61268/2kg/0806

Barcode

Batch No.:

Date of Manufacture:

® = Registered trademark of BASF

\* = Other trademarks